

Type Theory with General Variables (without the types) – after a proposal of Quine

Randall Holmes

March 9, 2025

0.1 Version Notes

3/9/2025: The proof that all formulas of TTGV are equivalent to stratified formulas is now I think fully written out, mod typos. Thomas, please read this carefully (pp 15-17 currently).

3/9/2025 7 pm: Some further revisions and corrections. Added the theorem about set abstracts.

Contents

0.1	Version Notes	1
1	Introduction	2
2	A Self Contained Development	2
2.1	Basic Axioms	2
2.2	Sets, atoms and individuals: some taxonomy	5
2.3	General set builder notation, the subset relation, power sets and unions	7
2.4	Relations, functions, and numbers	9
2.5	There are many kinds, but the apparent hierarchy of kinds is hard to talk about	13
2.6	Coercing formulas into typed or typable form	15

3	Typed theories of sets introduced; older proposals for type theory with general variables	20
3.1	Typed theories of sets: TST and variants	20
3.2	Quine’s original proposal of type theory with general variables	22
3.3	The system of Resnik	24
3.4	Ambiguity and stratification: NF and NFU	26
3.5	The proposal of Forster	27
3.6	Interpretation of the theories with general variables in the typed theories	28
4	Set theory without the axiom of pairing	31
5	More complex type theories	31

1 Introduction

This essay works on a proposal of Quine that type theory (the simple typed theory of sets, the precursor of New Foundations) can be presented as an unsorted theory.¹

In section 2, we give a self contained and naive presentation of a system of this kind from first principles. We call the system presented there TTGV (for “type theory with general variables”; Thomas Forster persists in referring to it as *trains de tres grand vitesse*.)

In section 3, we discuss prior proposals along these lines.

2 A Self Contained Development

2.1 Basic Axioms

In this document I will give a naive account from first principles of a theory of sets. It is ultimately based on an idea of Quine. We will talk about what this idea was in a later section when we shed our naivete. The theory is

¹This theory was defined by Holmes as a modification of Forster’s proposal which will be discussed later; at that time Holmes had no acquaintance with the earlier work of Quine or Resnik ([?]): it is a slight (?) weakening of Resnik’s theory, however, which confirms the naturalness of the approach.

called TTGV but we have no reason to use this name in this section, and explaining it would subvert our naivete.

The theory we introduce is a first order theory with membership and equality as primitive relations. We may in a less naive later part talk about the logic more carefully.

Some objects have elements and some do not. Objects that have elements are called *nonempty sets* (for the moment this is an unanalyzed phrase) and we state a natural identity criterion for nonempty sets.

Definition: An object is a nonempty set (written $\mathbf{Set}(x)$) if and only if it has an element:

$$\mathbf{Set}(x) \equiv_{\text{def}} (\exists y : y \in x)$$

Axiom of Extensionality: Nonempty sets with the same elements are equal:

$$(\forall xy : \mathbf{Set}(x) \wedge (\forall z : z \in x \leftrightarrow z \in y) \rightarrow x = y)$$

The axiom is not asymmetric between x and y : the hypotheses obviously imply $\mathbf{Set}(y)$.

The use of sets is to represent properties of objects: where P is a property of objects of kind κ , P is represented by the set of objects of kind κ which have property P , and we suppose that such a set exists for any kind and any property. We further suggest that objects belonging to the same set are of the same kind. Thus, objects are of the same kind if and only if there is a set which contains both of them. We view this as so important (for the sake of argument at least) that we adopt a definition and axioms.

Definition: Where x and y are objects, we define $x \sim y$, read, x is of a kind with y , as $(\exists z : x \in z \wedge y \in z)$.

Axiom of Kinds: For each object x there is an object $\kappa(x)$, which we call *the kind of x* , such that $x \in \kappa(x)$ and $(\forall y : y \in \kappa(x) \leftrightarrow y \sim x)$. Note that there is exactly one such object for each x by extensionality.²

²We refer to these sets as “kinds” rather than types, because a type is not really a set, but a syntactical characteristic of a variable. We do not extend this terminology to the historical systems discussed in section 3.

Theorem: The relation \sim is symmetric, reflexive³, and transitive [so the kinds are equivalence classes under this relation in some external sense].

Proof: $x \sim \kappa(x)$ and $x \sim \kappa(x)$, so $x \sim x$.

Logic tells us that if $x \sim y$, there is z such that $x \in z$ and $y \in z$, so $y \in z$ and $x \in z$, so $y \sim x$.

Suppose $x \sim y$ and $y \sim z$. Then $x \sim y$ and $z \sim y$. Then $x \in \kappa(y)$ and $z \in \kappa(y)$. So $x \sim z$.

This theorem motivates our giving ourselves permission to read $x \sim y$ as “ x and y are of the same kind”: they share a specific uniquely determined kind, they do not merely share some kind.

Now we talk about the construction of sets of objects of particular kinds with particular properties.

Axiom of Separation: For any object a and property $P(x)$ of objects x we introduce notation $\{x \in \kappa(a) : P(x)\}$ (this could also be written $\{x \sim a : P(x)\}$) and assert the following about the referents of these symbols:

1. For any b , $b \in \{x \in \kappa(a) : P(x)\}$ if and only if $b \sim a$ and $P(b)$.
This asserts the correspondence between properties and sets described in our motivation.
2. $\{x \in \kappa(a) : P(x)\} \sim \kappa(a)$
This asserts in effect that a kind is of the same kind as its subsets.
3. $\{x \in \kappa(a) : P(x)\} = \{x \in \kappa(b) : Q(x)\}$ iff $a \sim b$ (equivalently, $\kappa(a) = \kappa(b)$) and $(\forall x \in \kappa(a) : P(x) \leftrightarrow Q(x))$.

This provides identity conditions for the referents of these symbols. It only supplies new information here in the case where the extension of the referent of a set abstract is empty, but this is an important case.

Of course, this is an axiom scheme, with an instance for each formula $P(x)$ of our language defining a property.

³Forster notes that whether or not \sim is reflexive corresponds to whether we assume that everything belongs to something, that is, whether we are in set theory or class theory

Notice that $\{x \in \kappa(a) : P(x)\} \sim \kappa(a)$ is equivalent to

$$\{x \in \kappa(a) : P(x)\} \in \kappa(\kappa(a)).$$

Being of the same kind as a kind occurs often, and motivates a convenient definition.

Definition: We define $\kappa^1(x)$ as $\kappa(x)$ and define $\kappa^{n+1}(x)$ as $\kappa(\kappa^n(x))$ for each concrete numeral n (the n here is not a variable over which we can quantify).

Thus we can rewrite the observation above as

$$\{x \in \kappa(a) : P(x)\} \in \kappa^2(a).$$

Axiom of Diversity: If $\kappa(x) \sim \kappa(y)$ then $\kappa(x) = \kappa(y)$. No two distinct kinds are of the same kind. This is equivalent to the assertion that $\kappa^2(x) = \kappa^2(y) \rightarrow \kappa(x) = \kappa(y)$ [if you wonder why it is in the box with the definition above]. The purpose of this axiom will become clear below.

Definition: For any integer n , if a type $\kappa(u)$ belongs to $\kappa^n(x)$, define $\kappa^{n-1}(x)$ as $\kappa(u)$. If there is no such type, we leave this symbol undefined. This definition is justified by the Axiom of Diversity.

This completes the presentation of the basic axioms of the theory we are investigating. Unfolding the consequences will take a bit, since this is in fact a system which could be used as the foundation for mathematics.⁴ Further, its possible role as a foundation for mathematics suggests that we can expect that other candidate axioms will present themselves.

2.2 Sets, atoms and individuals: some taxonomy

We have introduced a notion of *nonempty set* at the beginning of our investigation. We now give a general account of sethood, which will allow us to investigate empty sets and non-sets of various kinds.

⁴The theory of Resnik, which we were not aware of when we defined this theory, differs from ours only in two additional assumptions which we can state at this point in terms of concepts we have explained: strong extensionality is assumed (objects with the same extension which cohabit with a nonempty set are equal), and an axiom provides that there are individuals (there is an object which does not cohabit with any nonempty set). Resnik apparently believed that he could prove that all individuals are of the same type, so this may be taken to be his intention, but it does not follow from his axioms.

Definition: We say that x is a set (written $\mathbf{set}(x)$) iff for some object y ,
 $x = \{z \in \kappa(y) : z \in x\}$.

Observation: It is straightforward to show by extensionality that if A is a nonempty set with $a \in A$ then $A = \{x \in \kappa(a) : x \in A\}$, since these will be nonempty sets with the same extension. So every nonempty set is a set.

Observation: Notice that every set belongs to a type $\kappa^2(y)$.

Definition: If x has the property that for no y is $\kappa(x) = \kappa^2(y)$ we call x an *individual*. Notice that no individual can have an element: if $u \in x$ then $x = \{v \in \kappa(u) : v \in x\}$ because these are nonempty sets with the same extension, and further $x \in \kappa^2(u)$.

Note further that there is nothing in our formalization which provides that two distinct individuals are necessarily of the same type, or that there are any individuals at all.

Definition: For any object x , we define $\emptyset_{\kappa(x)}$, the empty set cohabiting with $\kappa(x)$, or equivalently inhabiting $\kappa^2(x)$, as $\{y \in \kappa(x) : y \neq y\}$.

Clearly nothing can belong to $\emptyset_{\kappa(x)}$. Thus $\{y \in \kappa(x) : y \in \emptyset_{\kappa(x)}\} = \emptyset_{\kappa(x)}$, because the two are defined by set builder notations with the same bound and the same extension, so empty sets as we have defined them are in fact sets.

Observation: Note that $\emptyset_{\kappa(x)} \in \kappa^2(x)$. Note that any two empty sets of the same kind are equal: if u and v are sets of the same kind and nothing is an element of either, we have $u = \{x \in \kappa(w) : x \neq x\}$ and $v = \{x \in \kappa(q) : x \neq x\}$ and $\kappa^2(w) = \kappa^2(q)$, so $\kappa(w) = \kappa(q)$ by Diversity, so the two sets are equal because they have the same bounding kind and the same extension.

Nothing in this establishes that two elementless objects of the same kind are equal. This motivates the following definition.

Definition: An *atom* is an empty object which is not an individual (and so belongs to a kind which contains sets) but also not a set.

The theory does not prove the existence of either atoms or individuals but these are important formal possibilities.

Observation: Sets of the same kind with the same extension are equal. They are either nonempty sets with the same extension, so equal, or they are empty sets of the same kind, and so equal.

2.3 General set builder notation, the subset relation, power sets and unions

We define general set builder notation.

Definition: For any set $A \in \kappa^2(u)$ and property $P(x)$, we define

$$\{x \in A : P(x)\}$$

as $\{x \in \kappa(u) : x \in A \wedge P(x)\}$. The reader should be quite familiar with this notation: the only novel property to keep track of here is $\{x \in A : P(x)\} \sim A$ (which is directly verified).

We introduce the familiar subset relation, whose definition is a little more baroque than we are used to.

Definition: We define $x \subseteq y$ as

$$x \sim y \wedge \mathbf{set}(x) \wedge \mathbf{set}(y) \wedge (\forall z : z \in x \rightarrow z \in y).$$

Thorem and Definition: For any set A there is a set which we write $\mathcal{P}(A)$ such that for all B , $B \in \mathcal{P}(A)$ if and only if $B \subseteq A$. This set is called the power set of A .

Proof: It is evident that $A \subseteq A$ if A is a set. It is further evident that if $B \subseteq A$, then $B \sim A$. From this it follows that $\mathcal{P}(A)$ exists by the axiom of separation, being $\{B \in \kappa(A) : B \subseteq A\}$. Further, it is evident that $\mathcal{P}(A) \in \kappa^2(A)$.

Another very familiar notation is introduced.

Definition: If $x \sim y$, we define $\{x, y\}$ as $\{z \in \kappa(x) : z = x \vee z = y\}$. Note $\{x, y\} \in \kappa^2(x) = \kappa^2(y)$.

We define $\{x\}$ as $\{y \in \kappa(x) : y = x\}$. Of course this is the same as $\{x, x\}$ and belongs to $\kappa^2(x)$.

We define (x, y) as $\{\{x\}, \{x, y\}\}$ for any x, y of the same kind. Note that $(x, y) \in \kappa^3(x)$. Note that x can be determined from (x, y) as the only element which belongs to all elements of (x, y) , and y can be determined from (x, y) as the only element which belongs to exactly one element of (x, y) , from which it follows that if $(x, y) = (z, w)$, it follows that $x = z$ and $y = w$.

Observation: It is important to notice that we do not have an unrestricted axiom of pairing. The pair $\{x, y\}$ exists iff x and y are of the same kind.⁵ The same is true of ordered pairs, and this restricts the formation of relations and functions.

Definition: We define $\iota(x)$ as $\{x\}$. We define $\iota^0(x)$ as x and $\iota^{n+1}(x)$ as $\{\iota^n(x)\}$ for each concrete n (n here must be a numeral constant: no quantification is possible here). Notice that $\iota^n(x) \in \kappa^{n+1}(x)$ will hold for each concrete natural number.

We establish the existence of the union of a set.

Theorem and Definition: For any set A for which there is u such that $A \in \kappa^3(u)$, there is a set $\bigcup A \in \kappa^2(u)$ such that for all x , $x \in \bigcup A$ iff there is y such that $x \in y$ and $y \in A$.

Proof: The key is to show if $x \in y$ and $y \in A$, we can determine the type of x . We have $A \sim \kappa^2(u)$ and $A = \{u \in \kappa(x) : u \in A\} \sim \kappa(y)$, from which it follows by Diversity that $\kappa(y) = \kappa^2(u)$. We have $x \in y$ and $x \in \kappa(x)$, from which it follows that $y = \{u \in \kappa(x) : u \in y\}$ is equal to y and belongs to $\kappa^2(x)$. But then $\kappa^2(x) = \kappa^2(u)$, since both contain y , and it follows by Diversity that $\kappa(x) = \kappa(u)$, so by Separation we can define $\bigcup A$ as $\{x \in \kappa(u) : (\exists y : x \in y \wedge y \in A)\}$ and this belongs to $\kappa^2(u)$.

Definition: For $A \sim B$, sets, we define $A \cup B$ as $\bigcup\{A, B\}$, establishing the

⁵Notice that this is a restriction on applicability of the pairing operation, and TTGV falls in the more general category of Zermelo-like theories in which pairing fails. Track down Bolzano's discussion of attributes of the notion of set which includes formation of sets from objects of different species, which is not permitted here, and is permitted in Zermelo.

existence of binary unions⁶. We define $A \cap B$ as $\{x \in A : x \in B\}$ and define $A - B$ as $\{x \in A : x \notin B\}$. These operations are familiar though the fact that their operands must be of the same kind (and in all cases the resulting set is of the common kind) should be noted and verified.

We can now define $\{x_1, \dots, x_{n-1}, x_n\}$ as $\{x_1, \dots, x_{n-1}\} \cup \{x_n\}$, completing the definition of the usual notation for concrete finite sets by metatheoretical recursion.

2.4 Relations, functions, and numbers

We proceed to define relations and functions in a standard way, then define cardinal number in a way which is unusual now but has precedent.

Definition: Notice that if (a, b) exists, we must have a, b of the same kind and $(a, b) \in \kappa^3(a) = \kappa^3(b)$. Thus for any sets $A \sim B$ we can define $A \times B$ as $\{u \in \kappa^2(A) : (\exists ab : u = (a, b))\}$, and this set will contain exactly the pairs with first projection in A and second projection in B . As usual, this is called the cartesian product of A and B .

Definition: We call a set a *relation* if all of its elements are ordered pairs. If R is a relation, we define $x R y$ as $(x, y) \in R$. If R is a relation, we define $\text{dom}(R)$, the *domain* of R as $\{x \in \kappa^{-2}(R) : (\exists y : (x, y) \in R)\}$. We define R^{-1} as $\{u \in \kappa^0(R) : (\exists xy : x R y \wedge u = (y, x))\}$: this is called the *converse* of R . We define $\text{rng}(R)$, the *range* of R , as $\text{dom}(R^{-1})$. We define $R \text{``} A$ as $\text{rng}(R \cap (A \times \kappa^0(A)))$, for any set A and relation $R \in \kappa^3(A)$.

Definition: We call a relation F a *function* iff $F \text{``} \{a\}$ is a singleton set for each $a \in \text{dom}(F)$. We define $F(x)$ implicitly by the equation $F \text{``} \{a\} = \{F(a)\}$ for all $a \in \text{dom}(F)$. We define $F : A \rightarrow B$ as “ F is a function, $\text{dom}(F) = A$, and $\text{rng}(F) \subseteq B$ ”, and read this “ F is a function from A into B ”. We say that a function F is an *injection* iff F^{-1} is a function: if F^{-1} is a function, we call it the *inverse* of F . We say that a function

⁶It is amusing to note that the existence of binary unions immediately implies the axiom of diversity: if $\kappa(x) \sim \kappa(y)$ then $\kappa(x) \cup \kappa(y)$ exists and contains both x and y , so $x \sim y$ so $\kappa(x) = \kappa(y)$. In previous formulations we have used an axiom asserting the existence of binary unions of sets of the same type instead of the axiom of diversity. The reader can see above how diversity is used to prove set union (from which binary union follows) in this development.

is *onto* B iff $\text{rng}(F) = B$, and we say that a function is from A onto B if its domain is A and its range is B . We say that a function is a *bijection* from A to B if it is an injection with domain A and range B . We define $|A|$ as the set of all B such that there is a bijection from A to B , and call this the cardinality of A . The condition $|A| = |B|$ expresses for us the idea that two sets are of the same kind and the same size. There are obvious examples of sets which are of the same size but of different kinds: if $x \not\sim y$, then $\{x\}$ and $\{y\}$ are of the same size, but their cardinals are not the same, being of different kinds.

Definition: We define for any set A the set $\iota^{\text{“}}A$ as

$$\{u \in \kappa(A) : (\exists a \in A : u = \{a\})\}.$$

This is the elementwise image of A under the singleton operation. Notice that $\kappa(\iota^{\text{“}}A) = \kappa^2(A)$. We define $\iota^n^{\text{“}}A$ analogously. We define $T(|A|)$ as $|\iota^{\text{“}}A|$. It should be clear that this does not depend on the choice of A from the cardinal $|A|$. We define $T^n(|A|)$ as $|\iota^n^{\text{“}}A|$. An element of $|A|$ and an element of $T^n(|A|)$ are clearly of the same size, and this captures the relation of being the same size between sets of different kinds, but only under particular concrete circumstances. The T operation on cardinals is injective, so there are natural partial operations T^{-1} and T^{-n} .

Definition: We say that $|A| \leq |B|$ iff there is an injection from A to B (again, this clearly does not depend on the choice of representative sets from the cardinals). We define $|A| < |B|$ as holding iff $|A| \leq |B|$ and $|A| \neq |B|$.

Definition: We define $|A| + |B|$ as $|A \cup B|$ where A and B are disjoint. The sum of two cardinals will be undefined if they do not have disjoint representatives. We define $|A| \times |B|$ as $T^{-2}|A \times B|$. If A and B are sets, we define $|B|^{|A|}$ as $T^{-3}(|\{f : (f : A \rightarrow B)\}|)$, the cardinality of the set of functions from A to B shifted downward suitably in type. All of these operations may fail to be total. If the axiom of infinity is assumed, addition and multiplication of cardinals are total; exponentiation cannot be [because the collection of functions from A to B is provably larger than the common type of A and B under some circumstances].

Definition: We define $0_{\kappa^2(x)}$ as $|\emptyset_{\kappa(x)}|$ and $1_{\kappa^2(x)}$ as $|\{x\}|$. We say that a set of cardinals which belong to $\kappa^3(x)$ is *inductive* if it contains $0_{\kappa^2(x)}$ and contains $\lambda + 1_{\kappa^2(x)}$ whenever it contains λ . We define $\mathbb{N}_{\kappa^3(x)}$ as the intersection of all inductive sets with elements in $\kappa^3(x)$ [types containing sets of sets]. [the subscripting convention here is to subscript with a kind of the same kind as the subscripted object; the subscripts may be omitted where they can be understood in the context]. We have succeeded in defining natural numbers belonging to each kind $\kappa^3(x)$. Natural numbers of different kinds are not the same objects, but the T^n operation provides an injective external map from natural numbers in $\kappa^3(x)$ to natural numbers in $\kappa^{3+n}(x)$, which is also onto if the axiom of infinity is assumed.

Definition: We define a sequence as a function with domain some $\mathbb{N}_{\kappa^3(x)}$. We demonstrate the ability to define sequences by iteration such that $x_0 = a$ and $x_{i+1} = F(x_i)$. In this case we can write x_i as $F^i(a)$. We define the sequence x as the intersection of all sets which contain $(0, a)$ and contain $(u + 1, F(v))$ whenever they contain (u, v) : it is a straightforward exercise to prove that this set is a function with the desired properties. This can be used to justify more complicated recursive definitions. What it cannot do, except by analogy, is cast light on the meaning of notations $\kappa^n(x)$ or $\iota^n(x)$, because all elements of the domain of a sequence (or any function) must be of the same kind, and operations like κ and ι send objects to objects of different kinds.

Cantor's Theorem: We rearticulate and prove Cantor's Theorem on sizes of power sets. We have to rearticulate it because in its original form it is false: $|A| < |\mathcal{P}(A)|$ is false because the cardinals are of different kinds. The correct theorem is $|\iota^{\omega}A| < |\mathcal{P}(A)|$.

Proof of Cantor's Theorem: Clearly $|\iota^{\omega}A| \leq |\mathcal{P}(A)|$: the identity map is the desired injection.

Suppose $|\iota^{\omega}A| = |\mathcal{P}(A)|$. Then there is a bijection F from the singletons of elements of A to the subsets of A . All that we use about it here is that it is an injection. Define R as the set of all $a \in A$ such that $a \notin F(\{a\})$. Then consider $F^{-1}(R) = \{r\}$ and consider the proposition $r \in R$: this holds exactly if $r \notin F(\{r\}) = F(F^{-1}(R)) = R$, which is a contradiction.

The mathematics here is the same as in the usual treatment. The treatment of kinds here makes it a little less familiar.

Observation: Notice that for any x , $|\iota^{\kappa(x)}| < |\mathcal{P}(\kappa(x))|$: this indicates that there are more elements in $\kappa^2(x)$, which includes $\mathcal{P}(\kappa(x))$ as a subset, than there are (singletons of) elements of $\kappa(x)$. The cardinal $|\mathcal{P}(\kappa(x))|$ over $\kappa^2(x)$ is in an external sense larger than any cardinal over $\kappa(x)$. There is a proof here that $\kappa(x) \neq \kappa^2(x)$, though we do not address this issue until the next section. This is the treatment of the Cantor Paradox of the largest cardinal in this theory: there is a largest cardinal over each kind, but there are cardinals larger than this cardinal over the “next” kind.

Schroder-Bernstein Theorem: If $|A| \leq |B|$ and $|B| \leq |A|$ then $|A| = |B|$. The proof of this here is quite standard, we point out that we have it.

Observation: We can define transfinite ordinal numbers as isomorphism classes of well-orderings under similarity. The usual definition due to von Neumann does not work here, because the successor step $x \cup \{x\}$ of the construction of the usual ordinals always fails: there is no such set.

We define linear orders and well-orderings in the usual way: a well-ordering for us is reflexive (\leq rather than $<$). The notion of isomorphism of well-orderings is defined as usual. If \leq is a well-ordering, the order type of \leq , written $\text{ot}(\leq)$, is the isomorphism class of \leq ; an object is an ordinal number iff it is the order type of some well-ordering. For any relation R , we define R^ι as $\{(\{x\}, \{y\}) \in \tau^2(R) : x R y\}$ and for any ordinal $\alpha = \text{ot}(\leq)$ define $T(\alpha)$ as $\text{ot}(\leq^\iota)$. The ordinal $T(\alpha)$ seems in some sense to be the same order type as α , but it is a distinct object because it belongs to a different type.

Notice that if x is in the domain of a well-ordering \leq belonging to α , then \leq belongs to $\kappa^4(x)$ and α belongs to $\kappa^5(x)$. The well-ordering of the ordinals up to α , which we might think from the outside is the same order type as α , is actually in $\kappa^9(x)$, four types higher, and can be shown to be $T^4(\alpha)$. The Burali-Forti paradox does not afflict us: if Ω is the order type of the ordinals over $\kappa(x)$, the ordinal $\Omega + 1$ contains well-orderings longer (in an intuitive sense) than any over $\kappa(x)$, but it is a well-ordering over $\kappa^5(x)$ (the type where ordinals α over $\kappa(x)$

live) and we have here a proof that $\kappa^5(x)$ is not the same type as $\kappa(x)$, though we will not chase down the details (a simpler proof appears in the next section).

Observation: We can formulate axioms of infinity and choice very naturally. A simple form of the axiom of infinity is the assertion that for any natural number n (in any type) $n + 1$ is nonempty. The axiom of choice can be formalized exactly as usual as asserting that partitions have choice sets. The well-ordering theorem can be proved.

2.5 There are many kinds, but the apparent hierarchy of kinds is hard to talk about

We have an axiom, the Axiom of Diversity, which asserts that there cannot be more than one kind *of the same kind*. How do we know that there is more than one kind?

Theorem: For any x , $\kappa^2(x) \neq \kappa(x)$.

Proof: Suppose that we have an x such that $\kappa(x) = \kappa^2(x)$.

Define R as $\{y \in \kappa(x) : y \notin y\}$. $R \sim \kappa(x)$ so $R \in \kappa^2(x) = \kappa(x)$.

So $R \in R$ iff $R \in \kappa(x)$ [just shown to be true] and $R \notin R$. This is a contradiction.

Theorem: For each concrete $n > 1$, $\kappa^n(x) \neq \kappa(x)$.

Proof: The argument is very similar to the argument above but with some devious use of iteration of the singleton operation. Suppose $\kappa^n(x) = \kappa(x)$.

Let $R_{n,x} = \{\iota^{n-2}(y) \in \kappa^{n-1}(x) : \iota^{n-2}(y) \notin y\}$. To make it entirely clear that the existence of this set follows from Separation, rewrite it as $\{u \in \kappa^{n-1}(x) : (\exists y : u = \iota^{n-2}(y) \wedge \iota^{n-2}(y) \notin y)\}$. Notice that for any $y \in \kappa(x)$, $\iota^{n-2}(y) \in \kappa^{n-1}(x)$. Notice that $R_{n,x} \in \kappa^n(x) = \kappa(x)$. It follows that $\iota^{n-2}(R_{n,x}) \in \kappa^{n-1}(R_{n,x})$. It then follows that $\iota^{n-2}(R_{n,x}) \in R_{n,x}$ if and only if $\iota^{n-2}(R_{n,x}) \notin R_{n,x}$, which is impossible.

Observation: Another proof of this has already been given, perhaps not to be preferred because it is considerably more complicated. Cantor's

Theorem as stated and proved above, and the obvious fact that ι^A and ι^B are the same size iff A and B are the same size, imply that for any n , $|\iota^n(\kappa(x))| < |\iota^{n-1}\mathcal{P}(\kappa(x))| < \dots < |\iota^{n-i}\mathcal{P}^i(\kappa(x))| < \dots < |\mathcal{P}^n(\kappa(x))| \leq |\kappa^{n+1}(x)|$. Now if $\kappa(x) = \kappa^{n+1}(x)$ then there would be an obvious bijection from $\kappa(x)$ to $\iota^n(\kappa(x))$ (the singleton map, which would under this hypothesis be an actual function) and an obvious bijection from $\kappa(x)$ to $\kappa^{n+1}(x)$ (the identity map), contradicting this.

If we have an object x , we can now see that $\kappa(x)$, $\kappa^2(x)$, $\kappa^3(x)$ and so forth are distinct kinds, as far as we can count. But notice that we cannot really talk about this very much. We cannot define a sequence of kinds in this way, because when we define a sequence, all of its terms will be of the same kind.

Nor can we express the thought that this apparent sequence of kinds includes all the kinds there are. We just do not have the language to express this thought.

Now we can hint at another thing we cannot say. It seems very dubious to suppose that there is a kind $\kappa(x)$ such that $\kappa^i(x)$ is defined for every integer i . But we cannot prove that this is not the case, as we will discover. We may even be tempted to adopt an axiom which seems to imply that all kinds are like this.

We prove a lemma which we have in effect already proved, but which are useful to state here.

Typing Lemma: If $x \in y$ and $x \in \kappa^n(u)$, then $y \in \kappa^{n+1}(u)$. If $x \in y$ and $y \in \kappa^n(u)$, then $x \in \kappa^{n-1}(u)$.

Proof: Suppose $x \in y$ and $x \in \kappa^n(u)$. Any element of y is of the same kind as x and so belongs to $\kappa^n(u)$, so $y = \{v \in \kappa^n(u) : v \in y\}$, and this set belongs to $\kappa^{n+1}(u)$.

Suppose $x \in y$ and $y \in \kappa^n(u)$. It is also the case by the previous lemma that $y \in \kappa^2(x)$, so $\kappa^2(x) = \kappa^n(u)$, and $\kappa(x)$ belongs to $\kappa^2(x)$ and so is the unique type belonging to $\kappa^n(u)$, that is, $\kappa^{n-1}(u)$.

We prove a theorem with a famous name

Vicious Circle Principle: For each concrete n , there cannot be any sequence x_1, x_2, \dots, x_n such that each $x_i \in x_{i+1}$ and $x_n \in x_1$. That is, there are no loops (of a given concrete length) in the membership relation.

Proof: Suppose there was such a sequence. Clearly $x_i \in \kappa^i(x_i)$ for each concrete index i . Since $x_n \in x_1$, we have $x_1 \in \kappa^{n+1}(x_1)$. But by a theorem proved above, $\kappa^{n+1}(x_1)$ is distinct from $\kappa(x_1)$, which is also purported to contain $\kappa(x_1)$, and so disjoint from it (since kinds are equivalence classes under \sim) which is a contradiction.

2.6 Coercing formulas into typed or typable form

We discuss the possibility of imposing a typing convention on our language.

Definition: We say that a formula is *kind-bounded* iff each quantifier in the formula is restricted to a kind $\kappa(u)$, where u is free in the formula.

We say that a formula ϕ is *stratified* iff there is a function τ from variables to natural numbers such that for each subformula $x = y$ of ϕ we have $\tau(x) = \tau(y)$ and for each subformula $x \in y$ of ϕ we have $\tau(x) + 1 = \tau(y)$.

Meta-Theorem: Every kind-bounded formula in which we have an intended assignment of values to parameters is equivalent to a stratified formula.

Proof: We begin by defining a general procedure for assigning types in formulas which we highlight for future reference.

General typing procedure: Construct a function τ^* (in the metatheory, there can be no such function in our world) mapping variables in a formula ϕ to kinds. Each parameter is understood to be assigned a value, and since we know the value of a parameter v we can set $\tau^*(v) = \kappa(v)$. For each bound variable u which is restricted to $\kappa(u)$ where u is a parameter (whose value is understood to be known), we set $\tau^*(x) = \kappa(u)$. We can assume without loss of generality that bound variables restricted to different kinds have different names.

General stratification procedure: Now each atomic formula $u = v$ for which $\tau^*(u) \neq \tau^*(v)$ (both being defined) is equivalent to $\neg u = u$ and can be replaced with that, and each atomic formula $u \in v$ for which $\kappa(\tau^*(u)) \neq \tau^*(v)$ (both being defined) can be replaced with the equivalent $\neg u = u$.

In a kind-bounded formula ϕ , the general procedures above assign a type to every variable and make some modifications to subformulas to facilitate stratification enforcement.

In the formula ϕ^* obtained after these modifications, define $\tau(x)$ for each variable x as the largest n such that $\tau^*(x) = \kappa^n(\tau^*(y))$ for some variable y appearing in ϕ . This is a stratification.

Coercion to stratified form: We prove in this section that every formula of our language is equivalent to a stratified formula. We start (as above) with the stronger idea that each variable is assigned a kind as a type, in such a way that a parameter v will be assigned type $\kappa(v)$ and a bound variable x which is bounded in a kind $\kappa(u)$ will be assigned the same type as u .

Notice that if each of u and v is typed and $u = v$ appears in the formula and the types assigned to u and v are different, then $u = v$ can be replaced with $u \neq u$, since it is simply false (and this can be eliminated by logic). If $u \in v$ appears and u and v are both assigned types and the type assigned to v is not the kind of the type assigned to u , then $u \in v$ can be replaced with $u \neq u$ as it is false (and eliminated using logic). If the process of reduction removes all atomic subformulas under a quantifier, the entire quantified expression can be reduced to a truth value and eliminated. If all subformulas but one happen to be eliminated in the entire formula, surprisingly, the final formula can be a tautology or the negation thereof. Thus we can assume that these typing conditions hold on atomic subformulas in which both variables have been assigned types: for any atomic subformula $u R v$ where u is assigned type s and v is assigned type t , $s R t$ holds, where R is either $=$ or \in .

Notice that if we express each kind used as a type in the form $\kappa^n(u)$ where u is used as a type and is not the kind of any kind used as a type, then assigning n as type to each variable originally assigned $\kappa^n(u)$ will give a stratification [restricted to variables actually assigned type; a full stratification if all variables are typed] (subject to enforcement of the conditions on atomic subformulas described in the previous paragraph).

In a formula ψ in which each parameter and each variable bounded in the kind of a parameter are typed as discussed, consider a largest subformula $(\forall x : \phi)$ of ψ , where x is not assigned type because it is

quantified over without bound. Without loss of generality, we assume that we use only universal quantifiers, and that bound variables are systematically renamed so that different quantified subformulas always have different binding variables. The relation of *connectedness* between variables in a formula is defined as the smallest equivalence relation containing each pair of variables which appear together in an atomic subformula of the given formula.

In the formula ϕ , the variable x is free. Enumerate the types $\kappa(t_i)$ of the form $\kappa^m(u)$ where u is a type already used and m is an integer with absolute value less than or equal to the number of variables in ψ plus one. Notice that this will include all types which could be deduced from x by its occurrence with a variable already assigned a type in an atomic formula, but it also includes any type which could be deduced for x at any time in the future due to future type assignments to variables. $(\forall x : \phi)$ is equivalent to the conjunction of formulas $(\forall x_i \in \kappa(t_i) : \phi_i)$, where ϕ_i is the result of first replacing x with x_i then eliminating atomic subformulas which become ill-typed, and an exotic conjunct

$$(\forall x^* : (\bigwedge_i \kappa(x^*) \neq \kappa(t_i)) \rightarrow \phi^*),$$

where ϕ^* is the result of first replacing x with x^* , assigning it a type distinct from the $\kappa(t_i)$'s [the resulting formula ϕ^* will be the same for any such choice of type], then extending the type assignment and removing ill-typed atomic subformulas, which will include *all* of the atomic subformulas containing x and a variable already assigned a type, so that ϕ^* contains no typed free variables other than x^* which are connected to x^* 's via typed variables, and supports a type assignment to x^* , as well as the variables already typed [the only ones which occur are ones not connected to x^* , or at least not via variables assigned types]. Retain the type assignment to x^* for further use though we do not bound x^* in the formula. If x^* is connected to a variable y , there must be a variable z on the path from x^* to y which is untyped so far, which means it must be bound by an unrestricted quantifier, which must be inside ϕ^* since any quantifier in ψ whose scope includes $(\forall x : \phi)$ is restricted to a kind. [Further progress toward assigning a type to y will take place at the later stage where we process the formula quantified over z ; it is impossible for y to be assigned a type

used before the type of x^* was determined and remain connected to x^* when all variables on the path are typed, because the path cannot be long enough: some atomic subformula would be eliminated.]

Now we describe the elimination of the exotic conjuncts, whose strange boundedness to complements of concrete finite unions of kinds represent a stratification violation. [we write $\phi^*(x^*)$ instead of ϕ^* to facilitate writing many substitutions]

The assertion $(\forall x_1, \dots, x_{n+1} : (\bigwedge_{i=1}^{n+1} \bigwedge_{j=1}^{i-1} x_i \not\sim x_j) \rightarrow \bigvee_{i=1}^{n+1} \phi^*(x_i))$ succeeds in asserting that there are no more than n counterexamples $\kappa(t)$ to $(\forall x^* \in \kappa(t) : \phi^*(x^*))$ [in this transformation, note that a type assigned to x^* continues to be assigned to the variables x_i ; this is counterintuitive because the formula *says* that any two distinct x_i 's have different types, but formally it works, because we never do any type reasoning in contexts involving more than one of them subsequently, and in the formulas $x_i \not\sim x_j$, the two variables have the same relative type for purposes of stratification, weirdly enough].

Now it is straightforward to assert that listed items $\kappa(t'_1), \dots, \kappa(t'_m)$ are exactly the m counterexamples by asserting that each is a counterexample and there are no more than m counterexamples:

$$(\forall x_1, \dots, x_{m+1} : (\bigwedge_{i=1}^{m+1} \bigwedge_{j=1}^{i-1} x_i \not\sim x_j) \rightarrow \bigvee_{i=1}^{m+1} \phi^*(x_i)) \wedge \bigwedge_{i=1}^m \neg(\forall x'_i \in \kappa(t'_i) : \phi^*(x'_i))$$

Then the desired assertion equivalent to

$$(\forall x^* : (\bigwedge_i \kappa(x^*) \neq \kappa(t_i)) \rightarrow \phi^*(x^*))$$

is the disjunction of the assertions that each finite subset of $\{\kappa(t_1), \dots, \kappa(t_n)\}$ is the exact collection of counterexamples $\kappa(t)$ to $(\forall x^* \in \kappa(t) : \phi^*(x^*))$.

Now observe that this procedure produces a formula respecting all type assignments already made (preserving types assigned to unrestricted bound variables of which copies are made, as noted above).

We then iterate the procedure on the outermost quantifiers to which type assignments have not been made to the (unrestricted) binding variable, and we arrive ultimately at a completely typed formula, which

can then be stratified as described above [with the observation that subformulas $x_i \not\sim x_j$ produced in the elimination of exotic conjuncts are unproblematically stratified with x_i and x_j assigned the same type, as is easy to check by expanding them, though their semantics makes this rather unexpected.]

A further result which can be achieved by application of the methods of this section is that each instance of stratified comprehension

“ $\{x \in \kappa(u) : \phi(x)\}$ exists” is equivalent to an instance

“ $\{x \in \kappa(u) : \phi^*(x)\}$ exists” which is kind-bounded. It is straightforward to establish by purely logical manipulations that each instance is a consequence of ones in which each variable, free or bound, is connected to x : any subformula of a quantified formula in which no variable is connected to the binder can be pulled out of the quantified formula, and when all subformulas containing only variables (free or bound) not connected to x are pulled to the top, they can simply be replaced with truth values [each such replacement has the effect of making the original instance of comprehension a consequence of two simplified ones, one for each assignment of a truth value to the formula eliminated]. Now a formula ϕ in which each variable is connected to x will contain no non-trivial unbounded quantifiers at the end of the process described above, because the quantified variable is connected to x , and all the variables other than the quantified variable will have been assigned type, so *every* formula containing the quantified variable, and the quantifier itself, will be eliminated, so the stratified formula obtained at the end will be kind-bounded, and moreover every type assigned to a variable in it will be $\kappa^i(x)$ for some integer x .

It should be noted that all statements of equivalence here hinge on the instances of the separation axiom which prove $\kappa^n(x) \neq \kappa(x)$ for $n > 1$: these are used to justify elimination of ill-typed formulas.

3 Typed theories of sets introduced; older proposals for type theory with general variables

We presented TTGV in the previous section as if it were an independent proposal for the foundation of mathematics. The knowledgeable reader should be able to divine a lot about where it came from from what we have said so far; in this section we will make the historical background of this proposal clear. Our aim in organizing things this way is to make it clear that a theory of this kind can be presented without explicitly or implicitly supposing knowledge of the typed theories at all.

3.1 Typed theories of sets: TST and variants

The original theory of this kind, which appears to have been implicitly proposed by Norbert Wiener in 1914 and explicitly described by Tarski in the 1930s, has been called TST by the Belgian school of logicians who studied NF, and this is what we will call it. We note for historical accuracy that this is *not* the theory of types of Russell and Whitehead's *Principia*; it is considerably simpler, and Russell and Whitehead did not have mathematical knowledge required to simplify their system to this form.

TST is a multi-sorted first order theory with equality and membership. The sort of a variable v will be written $\mathbf{type}(v)$. We provide a countable supply of variables of each sort. Using $++$ to denote concatenation of strings, the formation rules for atomic formulas are that $v++' = '++w$ is a well formed atomic formula iff $\mathbf{type}(v) = \mathbf{type}(w)$, and $v++' \in '++w$ is a well formed atomic formula iff $\mathbf{type}(v) + 1 = \mathbf{type}(w)$. All atomic formulas are formed in this way. Writing this out in a way which manages use and mention correctly is a technical challenge!

We do not follow the convention of equipping variables with type superscripts in TST, which makes for very cluttered notation, though if we do provide a variable with a numeral superscript, one may expect that the type of that variable is as indicated.

The axioms of TST are a scheme of extensionality and a scheme of comprehension. The scheme of extensionality provides that each well-formed

formula of the shape

$$(\forall xy : x = y \leftrightarrow (\forall z : z \in x \leftrightarrow z \in y))$$

is an axiom. This asserts that objects of type $n + 1$ with the same extension (consisting of type n objects) are the same. The scheme of comprehension provides that for each well-formed formula ϕ in which the variable A is not free, $(\exists A : \forall x : (x \in A \leftrightarrow \phi))$ is an axiom if it is well-formed (the only additional requirement being that the type of A is the successor of the type of x).

It is usual to adjoin an axiom of infinity (whose form can be deduced from the development of mathematics in TTGV in the previous section) and often the axiom of choice to this theory, but they are not part of the formal definition of the theory we give here.

Some variants of this theory are worth noticing. Hao Wang proposed the variant TZT which differs simply by indexing the sorts by all integers instead of just the nonnegative integers. The consistency of TZT follows from the consistency of TST by a simple compactness argument. Wang himself called the theory TNT for “theory of negative types”; Forster prefers TZT because the theory in fact has all integer types, not just the negative ones.

The variant TSTU differs from TST in allowing urelements. Its extensionality scheme is

$$(\forall xyz : z \in x \rightarrow (x = y \leftrightarrow (\forall z : z \in x \leftrightarrow z \in y)))$$

(providing that nonempty sets with the same extension are equal) and it is convenient to supply a primitive constant \emptyset^{i+1} of each type i with the axiom scheme consisting of $(\forall x : x \notin \emptyset^{i+1})$ for each concrete natural number i . An object of type $i + 1$ is a set if it has elements or is equal to \emptyset^{i+1} .

In any of these theories, one can provide a term construction $\{x^i : \phi\}$ of a term of type $i + 1$ representing the unique set A such that $(\forall x : (x \in A \leftrightarrow \phi))$. The type rules for term constructions are straightforward to adapt from those for variables.

It is straightforward to show that TST is interpretable in the usual set theory ZFC. Let X_0 be an arbitrarily chosen set. Define X_{n+1} as $\mathcal{P}(X_n)$ for each n . In any formula of the language of TST, assign each parameter of type i a value in X_i and interpret each quantifier over type i as a quantifier restricted to X_i . It is straightforward to check that each interpretation of an

axiom of TST is true. The fact that the sets representing the types are not disjoint is harmless.⁷

We further note that TSTU is interpretable in TTGV, the theory we defined in the first section. Let $\kappa(x)$ be a kind. Interpret each parameter of type i as an element of $\kappa^{i+1}(x)$. Interpret each quantifier over type i as a quantifier restricted to $\kappa^{i+1}(x)$. That the axioms of TSTU hold is immediate from the axioms of TTGV: the weak extensionality of TSTU has the same form as the extensionality of TST, and the interpreted comprehension axiom of TSTU follows from the separation axiom of TTGV. Of course TST is interpretable in TTGV with the additional assumption of strong extensionality.

Our reasons for preferring to frame our flagship theory of types with general variables with weak extensionality will become evident shortly.

3.2 Quine’s original proposal of type theory with general variables

Quine’s original proposal of a type theory with general variables equivalent to TST is the subject of this subsection.

definition of “being of the previous type”: $x \text{ PT } y$ is defined as

$$(\exists zw : x \in w \wedge w \in z \wedge y \in z).$$

definition of type 0: $T_0(x)$ is defined as $(\forall y : \neg y \text{ PT } x)$.

definition of next type: For each concrete natural number n , $T_{n+1}(x)$ is defined as $(\forall y : T_n(y) \rightarrow y \text{ PT } x)$

He then stated his axioms schematically.

Quine’s comprehension axiom: For any formula ϕ ,

$$(\exists A : T_{n+1}(A) \wedge (x \in A \leftrightarrow (T_n(x) \wedge \phi))).$$

Quine’s extensionality axiom:

$$(\forall xyz : T_{n+1}(x) \wedge T_{n+1}(y) \wedge (\forall w : T_n(w) \rightarrow (w \in x \leftrightarrow w \in y)) \wedge x \in z \rightarrow y \in z).$$

I preserve the form of this axiom, which reflects defining equality in terms of membership, but it could be phrased differently.

⁷This is one of a couple of places in the paper where we are reminded that we need to track down Boffa’s discussion of typed properties

These are actually not the axioms as he first states them: this is the original extensionality axiom together with a modified version of the comprehension axiom which he states later as a consequence of the assumption that all elements of type $n + 1$ objects belong to type n , which his original axioms (astonishingly) do not imply.

Quine does recognize the importance of the relation of cohabitation as representing the notion of belonging to the same type, though he does not use it in the statement of his axioms.

This theory is not quite the same as ours. To begin with, it has what we regard as a formal defect: there is no need to axiomatize the theory with schemata with concrete natural numbers as indices, as we have demonstrated with our axiomatization (and as Resnik did prior to our work and very similarly). Quine does observe that he cannot prove and cannot even actually say that every object belongs to some type. Further, his theory says nothing at all about objects which do not belong to a type. In our theory and Resnik's, it is immediate that every object belongs to a type [which we call a "kind" in section 2], but the types may not be restricted to the familiar ones.

Quine says more about individuals than we do. Quine asserts that all individuals belong to the same type. Resnik also thought that he had asserted this (see below). We have not felt the need to do this, but we could. We also want to be free to explore the possibility there are no individuals at all.

We think that our presentation is superior to Quine's for a number of reasons. Our presentation does not allude to the simple typed theory of sets at all in its formulation [or much less obviously]: the fact that it is actually a presentation of the simple typed theory of sets unfolds in the development, as the reader should see in our first section. We dispute something that Quine says: he denies that systematic ambiguity as in Russell or in the development of New Foundations has a place here: in fact there is a strong place for systematic ambiguity in this theory; we do not escape this phenomenon when we transition to a one-sorted theory. But this also comes out in the development in section 2.

The axioms as selected above from Quine's treatment allow us to prove that all elements of a type $n + 1$ object are of type n : for any x of type $n + 1$ there is x^* of type $n + 1$ containing exactly the type n elements of x , and then by his original formulation of extensionality, $x^* = x$, so in fact all elements of x are of type n .

The fundamental point here is that Quine's theory is not intellectually independent from TST: Resnik's theory and mine are independent of TST(U)

in their formulation, though related notions naturally develop as these theories unfold.

Finally, our theory differs from Quine's quite deliberately in allowing atoms as well as empty sets, for reasons to be discussed soon.

NOTE: this system supports a proof that every formula is equivalent to the partial universal closure of a well-typed formula, because every type is associated with a formula, so the "all but finitely many types" restricted quantifiers can be handled. Details are different enough that they need to be written up. This discussion is needed because the comprehension axiom is stated for arbitrary formulas, which might suggest additional power over TST. In fact there is none.

3.3 The system of Resnik

What Quine did was a kludge. The presence of meta theoretic natural number parameters corresponding exactly to the types reveals that he is not really describing an autonomously motivated system.

Resnik gives a genuine one-sorted theory with one-sorted motivation from which type theory falls out, as we do, and his theory is very close to ours.

We list his seven axioms, staying closer to our own notation.

Definition: $x \sim y$ means $(\exists z : x \in z \wedge y \in z)$. Resnik defines $x = y$ as $(\forall z : x \in z \leftrightarrow y \in z)$. So does Quine; for us equality is a logical primitive, but the comprehension axiom of any of these theories should make this definition harmless.

Ax 1: $(\forall x : (\exists y : (\forall z : z \in y \leftrightarrow z \sim x)))$. This is almost the same as our axiom of types: ours has the extra clause $x \in y$ to ensure that \sim is reflexive. We use the notation $\tau(x)$ for the witness to the existential quantifier y . Strangely, the axiom of comprehension has to be used to fill in this detail in Resnik's system.

Ax 2: $(\forall xyw : y \in x \wedge y \in w \rightarrow x \sim w)$. Sets which meet have the same type. This is equivalent to our axiom of subsets. I'm wondering whether this is provable from Resnik's other axioms.

Ax 3: $(\forall uvwx : y \in x \wedge u \in x \wedge y \in w \wedge v \in w \rightarrow (\exists t : y \in t \wedge u \in t \wedge v \in t))$. This axiom is used to support transitivity of \sim . I believe that it is redundant. If $y \in x \wedge u \in x$ then we have $y \in \tau(u)$, where $\tau(u)$

witnesses Ax 1 with $x := u$. Similarly we have $v \in \tau(u)$. $u \in \tau(u)$ is not a consequence of Ax 1 (as it is in our formulation) but it does hold here because u belongs to some set by the hypotheses. So we can choose $\tau(u)$ as t .

Ax 4: $(\forall vwx yz : y \in x \wedge v \in w \wedge x \in z \wedge w \in z \rightarrow y \sim v)$ We say that in our development that because x and w have the same type, they have a union (a theorem following from the axiom of diversity), and of course this union will contain y and v . We have considered this exact statement as an axiom, but diversity seemed a simpler axiom.

Definition: $xPTy$ is defined (following Quine) as $(\exists zw : x \in w \wedge w \in z \wedge y \in z)$. $T_0(x)$ (x is an individual) is defined as meaning $\neg(\exists y : yPTx)$: nothing belonging to the same type as x has elements.

Ax 5: $(\exists x : T_0(x))$ We do not commit ourselves to the existence of any individuals. But it is natural to do so given the historical origin of this theory.

Ax 6: $(\neg T_0(x) \wedge x \sim y \wedge (\forall z : z \in x \leftrightarrow z \in y) \wedge x \in w) \rightarrow y \in w$. The form of this looks peculiar to us because Resnik treats equality is a defined notion, but it is the axiom of extensionality. It is a bit different from ours: it is weaker in that it does not force equality of nonempty sets with the same extension (Axiom 2 assists with this); it allows individuals with the same empty extension to be distinct but any empty object in a type containing a set is the only empty object in that type. This is natural; we are more liberal in allowing many atoms in each type.

Ax 7: For any formula ϕ , $(\forall z : (\exists y : w \sim z \wedge (\forall x : x \in y \leftrightarrow (\phi \wedge x \in z))))$. This is Zermelo's axiom scheme of separation, with the extra proviso that the set defined is of the same type as the bounding object (which actually doesn't have to be a set, and this has a use).

The maneuver for showing that a general object belongs to a set is rather strange here, and I want to be sure that Resnik actually realizes that he has to do it. For an arbitrary x , there is $w \sim x$ with empty extension, by axiom 7...and incidentally, some set contains both x and w , so x belongs to a set.

That said, this theory is the same as ours with stronger extensionality and the positive assertion that there are individuals. I think that my axiomatics

are cleaner, and that there are really good reasons to consider the possibility of atoms in addition to empty sets.

There is an error in Resnik. He claims that he can prove that all individuals are of the same type. This does not follow from his axioms. I think part of the issue is that he defines **ST** (the relation of being the same type, which we denote by \sim) in two different ways and does not seem to realize that they are not equivalent. It wouldn't be unreasonable given his evident intention to strength his axiom asserting the existence of individuals to assert as well that all individuals are of the same type.

We acknowledge this system as prior to ours, and as doing basically the same work: we were not aware of this work when we framed our system of the first section. We do think that there are formal advantages to our slightly (but inessentially) weaker system, which will come out in further discussion. Proofs of useful results from the first section port easily to this system.

3.4 Ambiguity and stratification: NF and NFU

TST exhibits a stronger form of a symmetry that Russell noted in the more complicated system of Principia Mathematica and called “systematic ambiguity”. This symmetry led to another proposal by Quine of an untyped version of TST which we describe because it is relevant to our project here.

In TST, provide a map ($x \mapsto x^+$) on variables which is an injection and raises type by one. For any formula ϕ , define ϕ^+ as the formula which results if each variable x is replaced with x^+ .

It is straightforward to see that if ϕ is provable, so is ϕ^+ . The converse is not true.

One could then reasonably conjecture the consistency of the Ambiguity Scheme, which asserts $\phi \leftrightarrow \phi^+$ for each closed formula ϕ .

Quine made the apparently stronger proposal that the types can simply be identified. The resulting theory is called NF (New Foundations) after the name of the paper in which it appeared.

NF is a one-sorted first order theory with equality and membership with the axiom of strong extensionality (objects with the same extension are equal) and the axiom scheme of stratified comprehension: $\{x : \phi\}$ exists if there is a function σ from variables to natural numbers such that each atomic subformula $v + + ' = ' + + w$ appearing in ϕ has $\sigma(v) = \sigma(w)$ and each atomic subformula $v + + ' \in ' + + w$ appearing in ϕ has $\sigma(v) + 1 = \sigma(w)$: such a formula is said to be stratified and the function σ is called a stratification.

Clearly it is equivalent to say that we are asserting that $\{x : \phi\}$ exists if ϕ could be turned into a well formed formula of TST by an appropriate assignment of sorts to variables.

NF presents difficulties: it was shown in 1953 to disprove Choice, and its consistency remained an open question until very recently.

Specker showed in 1962 that NF is equiconsistent with TST + Ambiguity, and with the existence of a model of TST in which there is a type raising endomorphism. This justifies Quine's jump from the temptation of the Ambiguity Scheme to the temptation of simply identifying the types.

Jensen showed in 1969 that NFU, the system with weak extensionality and stratified comprehension, is consistent and not even very strong. It is consistent with but does not prove Infinity, and it is consistent with Choice. This formal advantage of NFU over NF is the main reason that we choose to use weak extensionality in the definition of TTGV. It is now known that NF is consistent, but this is much harder to show, and the character of general models of NF remains ill understood.

3.5 The proposal of Forster

Thomas Forster proposed the following type theory of general variables, which was the first one I encountered. [NOTE: Thomas, I am lacking a copy of your initial communication with me about this: I surely have it but I cant find it in my email; this is a summary I put at the head of one of my versions].

This is a first order one sorted theory with equality and membership

Definition: $x \sim y$ is defined as $(\exists z : x \in z \wedge y \in z)$.

Axiom of weak extensionality: Objects with elements are equal if they have the same extension.

\sim is an equivalence relation: \sim is an equivalence relation in which the equivalence classes are sets: the equivalence class containing x is denoted by $\tau(x)$.

set union: The usual axiom of set union is asserted: for every A , $\bigcup A$ exists where $x \in \bigcup A \leftrightarrow (\exists y : x \in y \wedge y \in A)$

comprehension: $\{x \in \tau^n(u) : \phi\}$ exists where each variable appearing in ϕ is typed in the sense that (if it is a parameter) it belongs to some $\tau^m(u)$

and if it is bound, it is bound by a quantifier restricted to a $\tau^m(u)$, x is assigned type n of course, and further that in each subformula $u = v$ the types of u and v are the same and in each subformula $u \in v$, the type of v is the image under τ of the type of u .

This is something like the assertion that all sets determined by well typed set abstracts exist (using the terminology of the first section) though it is a bit less general.

The author is somewhat critical of this proposal of Forster, though it does have substantial interest. Like the proposal of Quine, it appears to depend philosophically on prior awareness of TST. It does have the interesting feature that it does not prove that the types are disjoint: if NF is consistent, a model of NF is a model of this theory in which there is only one type. Forster is also interested in the possibility of cycles in the types, in which $\tau^n(x)$ might be equal to $\tau(x)$ for some $n > 2$. These would correspond to type theories with loops in the types.

3.6 Interpretation of the theories with general variables in the typed theories

We now argue that a model of TST provides an interpretation of TTGV. These results will extend to the other theories, possibly under special assumptions.

Given a model of TSTU in which the sets implementing the types are disjoint (a model not satisfying this condition is readily modifiable to one which does), extend its language to a one-sorted language with the same variables by the device of assigning the value False to each ill typed atomic formula and interpreting complex formulas in the natural way.

All of the axioms of TTGV are obviously true in this structure for the language of TTGV except the axiom of separation. The problem with separation is that it asserts the existence of $\{x \in A : \phi\}$ for formulas which do not correspond to formulas for which this set is provided by the comprehension of TST.

We have foresightfully provided for this by proving in the first section that every set $\{x \in \tau(u) : \phi\}$ with fixed values for its parameters is provably equal to a set $\{x \in \tau(u) : \phi^*\}$ in which ϕ^* is well-typed, and in this context a well-typed formula is exactly equivalent to a well-formed formula of the underlying TSTU.

The system of Resnik is interpretable if the model of TSTU is a model of TST. This is direct, as Resnik's system differs very little from ours.

The system of Quine is close to TST in allowing only the types that TST itself has. Some application of a theorem similar to our well-typedness theorem will be needed, because the comprehension axiom of Quine's system is not restricted to well-typed formulas.

The system of Forster shines here, because its comprehension axiom provides for exactly the sets which the comprehension scheme of TST provides for, and also there is no need for the condition that the types are disjoint which is important in our proof above.

Quine's system is in some sense exactly equivalent to TST (apart from the possibility of objects not satisfying any of the predicates T_n , about which the theory says nothing).

The other theories cannot be said to be exactly equivalent to TST, because they do not restrict themselves to the hierarchy of types indexed by the natural numbers which TST supports, and in fact their language cannot even express such a restriction.

We describe a theory TSTG which is typed and in some sense exactly equivalent to TTGV.

TSTG is a first order multisorted theory (or family of theories) with sorts of two kinds, $\tau^+(l, n)$ where l is a label and n is a natural number, and $\tau(l, i)$ where l is a label and i is an integer (a version of the theory might have only one of these kinds of type). For any type t we define t^+ as $\tau^+(l, n + 1)$ if $t = \tau^+(l, n)$ and as $\tau(l, i + 1)$ if $t = \tau(l, i)$.

An atomic subformula $u = v$ is well-formed ff the types of u and v are the same; $u \in v$ in which u is of type t is well-formed if and only if the type of v is t^+ .

The weak extensionality and comprehension axioms of TSTG are the complete schemes of formulas of the same shapes given for TSTU, with the additional latitude afforded by having more types.

In effect, we are providing for an arbitrary large collection of models of TST and an arbitrarily large collection of models of TST. This is a family of theories because we have not stipulated how many labels there are for types of each kind.

Now there is a direct translation between models of TTGV and models of TSTG. From a model of TTGV obtain a model of TSTG in which the sets implementing the types of TTGV are the extensions of the kinds $\kappa(x)$ in the model of TTGV, and the $(t \mapsto t^+)$ operation on type labels parallels the

κ operation on types in the sense of TTGV (to realize the type labels, one needs to make a choice of “base type” in each orbit in κ without a minimal element; this is not an essential use of choice because we could also allow many interconvertible notations for each type in a sequence of types indexed by all integers). This is readily seen to be a model of TSTG.

A model of TSTG is converted to a model of TTGV by assigning values to all ill-typed atomic formulas of False (ensuring first that the sets implementing the types are pairwise disjoint) and extending the definition of truth values of general formulas appropriately. Again, the only axiom of TTGV which requires care to verify in the resulting structure is separation, and its validity follows from the fact that general set abstracts in TTGV are equivalent to well-typed set abstracts.

This result adapts to the theory of Resnik which has the added assumptions of strong extensionality and existence of individuals.

Consideration of the theory TSTG can be useful in thinking out things about TTGV. Notice that an arbitrary set of models of TSTG can easily be made a pairwise disjoint set, and the union of a pairwise disjoint collection of models of TSTG is a model of TSTG. TTGV has similar properties.

One way to get the condition that any well-typed formula $\phi(\kappa(y)/x)$ in which x is the only free variable in ϕ will hold for all values of $\kappa(y)$ if it holds for all but a concrete finite collection of values of $\kappa(y)$, which is sufficient to make every formula equivalent to the partial universal closure of a well-typed formula, is to replace the model of TTGV in which one works with a countable union of pairwise disjoint copies of the model one starts with.

Another observation is that we cannot (verifying a claim we made above) establish that arithmetic is the same everywhere in a model of TTGV, because we can take unions of models of TST with different arithmetic facts and convert them to a model of TTGV.

Note that consistency of NFU (and of NF) implies consistency of TTGV (even with strong extensionality) with the Ambiguity Scheme which asserts that for any formula ϕ in which x is the only free variable,

$$(\forall uv : \phi[\kappa(u)/x] \leftrightarrow \phi[\kappa(x)/x]).$$

A model of TSTU with a type shifting endomorphism, which exists by the results of Jensen and Specker, converts to a model of TTGV in which this is true.

An interesting footnote to this section is that TTGV is not finitely axiomatizable. The separation axiom for typed formulas is finitely axiomatizable:

this can be done for example by converting the axioms of Hailperin's finite axiomatization of NF to well-typed formulas closed with a quantifier over all types. The equivalence of set abstracts over general formulas to set abstracts over well-typed formulas depends also on the disjointness of the types, which depends on a countable collection of instances of TTGV comprehension (the ones defining the sets $R_{n,x}$ in the proof). Suppose that TTGV was finitely axiomatizable. Each of the axioms in this finite axiomatization would be provable using the axioms implementing well-typed separation and finitely many of the axioms providing for $R_{n,x}$'s (and the other axioms of TTGV). Thus there would be a finite axiomatization consisting of the well-typed separation axioms, the other axioms of TTGV and finitely many of the axioms providing $R_{n,x}$'s. But all of these axioms hold in a model of cyclic type theory with weak extensionality with a large enough finite number of axioms: because NFU is consistent, this theory is consistent. And this theory does not cover TTGV because it does not prove existence of one of the $R_{n,x}$'s.

4 Set theory without the axiom of pairing

5 More complex type theories

A weird feature of the theory TTGV is that it is impossible to establish any relation between types where neither type is an image under an iterate of κ of the other.

This can be amended by the maneuver of providing the general ordered pair (x, y) as a primitive construction, with the following axioms:

Axiom of ordered pairs: For any x, y, z, w , $(x, y) \sim (z, w)$ if and only if $x \sim z$ and $y \sim w$, and $(x, y) = (z, w)$ iff $x = z$ and $y = w$.

These axioms are sufficient to provide the set $A \times B = \{(a, b) : a \in A \wedge b \in B\}$, the cartesian product, and for any set of ordered pairs R the inverse $R^{-1} = \{(b, a) : (a, b) \in R\}$ and $\text{dom}(R) = \{a : (\exists b : (a, b) \in R)\}$.

This has the effect that cardinalities of sets of diverse kinds can be compared, which can for example be used to prove that there are types of individuals. In any given type, one can define the collection of sets which are of the same cardinality as some type. Observe that the minimal ordinal not the order type of a well-ordering of a subset of $\kappa(x)$ must be less than the

minimal ordinal not the order type of a well-ordering of a subset of $\kappa^2(x)$; this fact could not even be expressed without the axiom of pairs. Consider types $\kappa(x)$ for which the minimal order type of a well-ordering of a subset of $\kappa(x)$ takes the smallest possible value (for types whose cardinality can be seen in a particular type). This must be a type of individuals.

This axiom allows one to interpret the type system of Principia which had arbitrary relation types with heterogeneously typed arguments.

Further, with a bit of ingenuity, one can present the modern theories of dependent types in this way.

The quick and dirty way to do this is first to add impredicative class theory on top of TTGV with pairing. Impredicative class theory by itself allows the proof that there are types of individuals by a method resembling the one just given.

Then add dependent pairs: for every class function f with a type as its domain, construct dependent pairs (x, y, f) where $y \in f(x)$, and declare that $(x, y, f) \sim (u, v, g)$ if and only if $x \sim u$, $y \sim v$, and $f = g$, while $(x, y, f) = (u, v, g)$ iff $x = u$, $y = v$, and $f = g$.

This gives a quite strong dependent type theory with a single universe.

A universe would be a class which is the domain of a class function whose range is closed under desired type constructions. We will articulate this in more detail when we have thought out the details.

Finally, it might be possible to remove the impredicative character of the class theory by exploiting the universes cleverly (and defining them with care).

Another stab at this:

Start by adding the predicative theory of classes (class comprehension has all quantifiers bounded in classes).

Then add the pair – for any two objects x, y there is a pair (x, y) and $(x, y) = (z, w)$ implies $x = z$ and $y = w$, and $x \sim z$ and $y \sim w$ implies $(x, y) \sim (z, w)$ [notice this is not a biconditional: we define relations not as general sets of ordered pairs but as subsets of Cartesian products, and this is enough to get domains and inverses)].

The absolute pair enables comparison of cardinality of sets of different types. $|A| = |B|$ is defined to mean “there is a bijection between A and B ”.

Now we adopt a weak axiom of univalence: for each nonempty set a , there is exactly one type $\pi[a]$ such that $|a| = |\pi[a]|$ and moreover we provide a bijection $\Pi_a : \pi[a] \rightarrow a$.

This is philosophically amusing as a strong assertion of data type security.

It also has an unintended consequence in this particular theory: for any cardinals κ and λ , if $2^\kappa = 2^\lambda$ then $\kappa = \lambda$.

It means that we can define type constructors in quite standard ways. $\pi[\pi(x) \times \pi(y)]$ is a type in which the elements are exactly correlated with [pairs of an object of the type of x and an object of the type of y . There is no guarantee that this will hold of the set $\pi(x) \times \pi(y)$ of absolute pairs: every such pair will be in this set, but there may be other objects in the same type as the elements of $\pi(x) \times \pi(y)$.

Now we say that a class V is a universe iff $x \in V \rightarrow \pi(x) \in V$ for all x and for any class function f with domain a nonempty set $A \in V$ and all range elements nonempty and belonging to V there is an entity $\lambda_f \in V$ such that all $((a, b), \lambda_f)$ with $a \in A$ and $b \in f(a)$ are of the same type. This is a little different from what I did above, since I use the same absolute pair here.

We then assert as an axiom that every object belongs to a universe.